



TITLE:

Recovery of genetic diversity in threatened plants through use of germinated seeds from herbarium specimens

AUTHOR(S):

Nakahama, Naoyuki; Hirasawa, Yuki; Minato, Tsubasa; Hasegawa, Masahiro; Isagi, Yuji; Shiga, Takashi

CITATION:

Nakahama, Naoyuki ...[et al]. Recovery of genetic diversity in threatened plants through use of germinated seeds from herbarium specimens. *Plant Ecology* 2015, 216(12): 1635-1647

ISSUE DATE:

2015-12

URL:

<http://hdl.handle.net/2433/207704>

RIGHT:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s11258-015-0547-8>; The full-text file will be made open to the public on 1 December 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Recovery of genetic diversity in threatened plants through use of germinated seeds from herbarium specimens

Naoyuki NAKAHAMA¹, Yuki HIRASAWA², Tsubasa MINATO³, Masahiro HASEGAWA⁴, Yuji ISAGI¹,
Takashi SHIGA^{2,3}

¹ Graduate School of Agriculture, Kyoto University, Oiwake-cho, Kitashirakawa, Sakyo-ku,

Kyoto 606-8502, Japan

² Graduate School of Education, Niigata University, 8050, Ikarashi2-no-cho, Nishi-ku, Niigata

950-2181, Japan

³ Faculty of Education, Niigata University, 8050, Ikarashi2-no-cho, Nishi-ku, Niigata 950-

2181, Japan

⁴ Osaka Museum of Natural History. Nagai Park 1-23, Higashisumiyoshi-ku, Osaka 546-0034,

Japan

Corresponding author: Naoyuki Nakahama

E-mail: nakahama.naoyuki.68r@st.kyoto-u.ac.jp,

The Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku

Kyoto 606-8502, Japan.

Tel: +81 75 753 6129, Fax: 81 75 753 6129

5003 words (Abstract 195 words)

Abstract

The reintroduction of *ex situ* conserved individuals is an important approach for conserving threatened plants and reducing extinction risk. In this study, we elucidated the effects on the genetic diversity of wild populations of *Vincetoxicum pycnostelma* Kitag. [= *Cynanchum paniculatum* (Bunge) Kitag.] by modelling the genetic consequences of reintroducing plants using the germinated seeds of herbarium specimens. This semi-natural grassland herb is threatened in Japan. First, we tested the germinability of seeds from herbarium specimens collected from nine sites in Kinki and Tokai districts, Japan (one specimen per site, total 206 seeds). Next, we analysed the genetic diversity and structure of germinated seedlings and the current wild individuals using nine polymorphic microsatellite markers. Germination was observed for 38 seeds (18.4%) from four specimens collected 3–18 years prior to the study. Although the genetic diversity of the specimens' seeds was lower than that of the wild population because of the small sample size, the seedlings from specimens taken from three sites had unique alleles that did not exist in the wild populations. Consequently, viable herbarium specimens' seeds with unique alleles could be useful resources for recovering the genetic diversity in threatened wild plant populations.

Keyword: threatened plants, genetic diversity recovery, germination test, microsatellite, reintroduction

Introduction

Many species worldwide face an extinction crisis because of the destruction and fragmentation of habitats as a result of human activities (Ceballos and Ehrlich 2002; Hooper et al. 2005; Aguilar et al. 2006). To conserve the populations of threatened species, it is crucial to maintain not only population size and habitat but also genetic diversity for preventing the decline in reproductive success owing to inbreeding depression, which would escalate the extinction risk (Keller and Waller 2002; Mattila et al. 2012; Palomares et al. 2012). Furthermore, a loss of genetic diversity would compromise their adaptive potential, particularly in the face of concerns pertaining to climate change (Blows and Hoffmann 2005; Willi et al. 2006).

Although the conservation of natural habitat is critical for conserving ecological interactions, *in situ* and *ex situ* conservation is also appropriate for safeguarding individual species against extinction in the wild. For example, use of soil seed bank is an *in situ* natural resource for the reproduction of plants that is effective for the restoration of genetic diversity (Uesugi et al. 2007; Zaghoul et al. 2013). On the other hand, the primary purpose of *ex situ* conservation is to maintain wild species outside their natural habitat so that species recovery and reintroduction can be attempted if the wild populations severely decline or become extinct. It is well known that preserved seeds are useful as *ex situ* populations (Honney et al. 2008; Frankham et al. 2009; Hoban and Schlarbaum 2014). The reintroduction of many individuals germinated from preserved seeds also allows the recovery of population size and genetic diversity (Honney et al. 2008; Guerrant et al. 2014; Hoban and Schlarbaum 2014). Thus, projects to preserve the seeds of many plants are used worldwide and comprise an efficient method for plant reintroduction and conservation (Schoen and Brown 2001; Guerrant et al. 2014; Hoban and Schlarbaum 2014). For example, the ‘Millennium

64 Seed Bank Project' of the Royal Botanic Gardens, Kew, the 'Svalbard Global Seed Vault' and many
65 other botanical gardens store wild and cultivated plant seeds as a resource for future use in
66 conservation translocations and crop development (Schoen and Brown 2001; Van Slageren 2003;
67 Qvenild 2008; Alsos et al. 2013).

68 However, re-establishment of the seeds collected in the past has risks pertaining to inbreeding
69 and inbreeding depression if the sources are restricted or population size is small (Frankel and Soulé
70 1981; Schoen and Brown 2001). In addition, reintroduced individuals collected from remote habitats
71 would also have reduced fitness because of local adaptation (Becker et al. 2006; Leimu and Fischer
72 2008; Hereford 2009). Furthermore, reintroduced individuals that are genetically distinct from a wild
73 population would not only have a risk of outbreeding depression (Montalvo and Ellstrand 2001; Huff
74 et al. 2011), but would also lead to the loss of the local genetic identity of the native population
75 (Gottelli et al. 1994; Milián-García et al. 2014). Accordingly, to conserve threatened plants by
76 reintroducing individuals from preserved seeds, it is very important to collect many seeds from
77 widely dispersed populations (Falk and Holsinger 1991; Hoban and Schlarbaum 2014). However,
78 collecting seed resources of threatened or locally extinct species from wild populations may be
79 difficult, although seed banking projects throughout the world have been recently constructed and
80 followed the protocols to ensure the maintenance of genetic diversity (León-Lobos et al. 2012;
81 Guerrant et al. 2014; Hoban and Schlarbaum, 2014). Herbarium specimens in museums can retain
82 viable and germinable diaspores (Windham et al. 1986; Bowles et al. 1993; Lledó et al. 1996;
83 Magrini et al. 2010; Magrini 2011; Bewley et al. 2013; Shiga 2013); therefore, they have potential as
84 a reintroduction resource. Furthermore, the viable diaspores in herbaria specimens are also useful for
85 the conservation of threatened local populations in each plant species because presumably the

specimens are collected from these populations. Despite the potential use of herbarium specimens' seeds, the effect of reintroduced seeds on the restoration of genetic diversity has not been studied, and it is unknown whether genetic diversity now lost in wild populations can be found in herbarium collections.

Vincetoxicum pycnostelma Kitag. [= *Cynanchum paniculatum* (Bunge) Kitag.] (Fig. 1a), subfamily Asclepiadoideae of Apocynaceae, is a perennial herb that grows in semi-natural grasslands in Japan, Korea, China, Mongolia and Russia (Yamazaki 1993; Wu and Raven 1995; Flora of Korea Editorial Committee 2007). Although this species was common a few decades ago, its populations have rapidly declined because of land-use changes in semi-natural grasslands in Japan (Environment Agency of Japan 2000; Uematsu et al. 2010). It is estimated that the extinction probability after 100 years is 96% (Environment Agency of Japan 2000). The species is categorized as 'near threatened' in the Japanese Red List (Ministry of the Environment Government of Japan 2012). Many other native herbaceous plants have also experienced a rapid decline in semi-natural grasslands (Koyanagi and Furukawa 2013); hence *V. pycnostelma* is a prime example of these rare and threatened species.

In this study, we examined the germinability of seeds from herbarium specimens of *V. pycnostelma* and we assessed the effect on the genetic diversity and structure of the wild populations by modelling the reintroduction of germinated seeds. We also discuss suitable methods for collecting and managing herbarium specimens' seeds at museums that are to be used for conserving not only threatened plants but also the local populations in each species.

Materials and methods

Selection of herbarium specimens

Seeds were collected from the herbarium of the Osaka Museum of Natural History, Japan (Figs. 1a, b). The museum was established in 1974, and the herbarium preserves more than 270,000 vascular plant specimens (Osaka Museum of Natural History 2012). In this herbarium, insect control involved naphthalene application and fumigation with carbon disulphide; the room temperature was not controlled until 2000. However, since 2001, integrated insect control has been performed by applying naphthalene and freezing treatments and the room conditions have been controlled at 20°C and 50% humidity. In the herbarium, 206 seeds from nine specimens with mature fruits (one specimen per site) from Kinki and Tokai districts, Japan, had been collected (Fig. 2). The ages of the nine stored specimens were from 4 to 62 years (Table 1). Seeds were collected from only one fruit per specimen. After the germination test, specimens of the seedlings or seeds were mounted and donated to the Osaka Museum of Natural History with annotation cards.

Germination test of herbarium specimens' seeds

Germination of the herbarium specimens' seeds was tested using the screening test system (Washitani 1987) under a 12/12-h photoperiod and two temperature regimes: 1) a gradually increasing temperature regime from 4°C to 36°C at intervals of 4°C (IT) and 2) a gradually decreasing temperature regime from 36°C to 4°C at intervals of 4°C (DT). The two treatments were conducted because the preservation condition in the herbarium may have affected the dormancy of the specimens' seeds, and it was predicted that a cold stratification treatment would be required to break the seed dormancy of *V. pycnotelma* (Zhou et al. 2003). Considering that growth rate increases with temperature, the duration of the higher temperature treatment was less than the duration of the lower temperature treatment (Table S1). After reaching the final temperature in each regime, these

seeds were incubated at 25°C (DT, 5 days) or treated at alternating temperatures of 12°C and 25°C at 24-h intervals (IT, total 5 days). The number of germinated seedlings was counted when the temperatures were changed, and the seedlings were moved to an incubator at 25°C to determine whether they would grow normally. For microsatellite analysis, we selected well-growing plants among 32 samples of 38 seedlings obtained from four specimen sheets. The germination test was conducted in June 2012 (Site i) and May 2013 (Sites a–h; Table 1).

The ungerminated seeds were tested using tetrazolium (2,3,5-triphenyl tetrazolium chloride) to examine their viability (Cottrell 1947; Elias et al. 2012). The seeds were cut and stained for 48 h in the dark with 1% tetrazolium solution, and seeds that stained red were defined as viable.

Sampling and microsatellite analysis

In 2012 and 2013, leaf samples from 131 adult individuals were collected from sites same as those from where the herbarium specimens had been collected (Figs. 1c and 2). The samples were used to estimate the genetic diversity among the extant populations (Table 2). At all sites, we comprehensively collected samples from each entire patch. Because *V. pycnotelma* at Site e was locally extinct, we collected leaf samples from a neighbouring population located one kilometre away. The number of individuals was counted at each site.

Genomic DNA was extracted using a modified cetyltrimethylammonium bromide method (Milligan 1992). The genotypes of each individual, including wild populations and specimens' seedlings, were characterized at nine microsatellite loci. Seven of the nine loci were characterized by Nakahama et al. (2012): *Vpy002*, *Vpy006*, *Vpy012*, *Vpy013*, *Vpy16*, *Vpy018* and *Vpy022*. Two of the nine loci were developed by Nakahama et al. (unpublished data): *Vpy025* (GenBank accession

number: AB948217) and *Vpy031* (GenBank accession number: AB948218). Polymerase chain reaction (PCR) amplifications, except for those of *Vpy025* and *Vpy031*, were performed following the standard protocol of the Qiagen Multiplex PCR kit (Qiagen) with a final reaction volume of 5 μ L, which contained 16 ng extracted DNA, 2.5 μ L of 2 \times Multiplex Master Mix and 0.2 μ M of each multiplexed primer. For *Vpy025* and *Vpy031*, the forward primer was synthesized with the M13 tag sequence (*Vpy025* 5'-CACGACGTTGTAAAACGAC-3', *Vpy031* 5'-TGTGGAATTGTGAGCGG-3'; Boutin–Ganache et al. 2001). The PCR mixtures of *Vpy025* and *Vpy031* had a final volume of 5 μ L, which included 16 ng extracted DNA, 2.5 μ L of Multiplex PCR Master Mix, 0.01 μ M forward primer, 0.2 μ M reverse primer and 0.1 μ M M13 fluorescent primer. The PCR amplifications of all loci were carried out using a GeneAmp PCR System 2700 thermal cycler (Applied Biosystems, Tokyo, Japan) using the following conditions: initial denaturation at 95°C for 15 min, followed by 25 cycles of 30 s at 94°C, 1.5 min at 57°C and 1 min at 72°C, and a final extension for 30 min at 60°C. The PCR product size was measured using an ABI PRISM 3130 Genetic Analyzer (Applied Biosystems) and GeneMapper ver. 4.1 (Applied Biosystems).

Statistical analysis of the genetic diversity and structure

Genetic diversity was evaluated for the specimens' seedlings, the wild populations and the hypothetical mixed populations, which comprised a hypothetical mixing of the specimens' seedlings and each corresponding wild population. The genetic diversity was evaluated in terms of the following: average number of alleles per locus, allelic richness (El Mousadik and Petit 1996), summed number of rare alleles with frequencies less than 5% among the total population, expected heterozygosity, observed heterozygosity and inbreeding coefficient. We also evaluated the summed

number of unique alleles that were only present in specimens' seedlings from each site. All of these parameters, except allelic richness summed number of rare and unique alleles, were calculated using GenAlEx version 6.41 (Peakall and Smouse 2006). Allelic richness, deviation from Hardy–Weinberg equilibrium and the linkage disequilibrium between loci were also determined using FSTAT ver. 2.9.3 software (Goudet 2001). The significances of the heterozygosity excess and deficit values were tested by comparison with the 95% confidence intervals derived from 108,000 randomizations. We evaluated the genetic relationships between the wild populations at all sites using Bayesian clustering with STRUCTURE ver. 2.3.4. (Pritchard et al. 2010), which assigns individuals into K clusters. The population structure was simulated with the values of $K = 1–8$ under an admixture model and the correlated allele frequency model (Hubisz et al. 2009). All runs involved one million Markov chain Monte Carlo iterations after a burn-in period of one million iterations. Twenty runs were performed for each value of K . The F value, the estimated amount of genetic drift between each cluster and a common ancestral population and the expected heterozygosity were calculated. The number of clusters was determined by comparing the mean values and the variability of log likelihoods for each run. To select the optimal value of K , STRUCTURE HARVESTER was used (Earl and vonHoldt 2012). We also evaluated genetic relationships between the wild populations and specimens' seedlings at each site by the same method.

To evaluate genetic differentiation between specimens' seedlings and wild populations for each site at the individual level, we calculated pairwise co-dominant genotypic distances (Smouse and Peakall 1999) between all specimens' seedlings and wild individuals from all sites. We also performed principal co-ordinates analysis (PCoA) using GenAlEx version 6.41 (Peakall and Smouse 2006).

Results

Germination and viability of herbarium specimens' seeds

A total of 38 seeds, representing four herbarium specimens, were germinated out of a total of 206 seeds (Fig. 1d and Table 1). The germination percentage for seeds from each specimen ranged from 0% to 56.5%, with an average of 18.4%. The oldest germinated specimens' seeds were collected 18 years prior to the study. Under the IT and DT conditions, seed germination from two and four specimens were observed, respectively. Each 18.4% of the seeds germinated in both conditions. Furthermore, 3.0% of ungerminated seeds were confirmed to be viable according to the tetrazolium dye test (Table 1). The viable seeds were collected from three herbarium specimens, which were collected 9 to 18 years prior to the study. The ungerminated viable seeds were confirmed only in the IT condition.

Genetic diversity of seedlings from herbarium specimens, wild populations and hypothetical mixed populations

There was no evidence for large allele dropouts or null alleles in the data set. The numbers of alleles, the allelic richness and the expected heterozygosity of herbarium specimens' seedlings was considerably lower than those of the wild populations because their allele frequencies were dominated by a few alleles, although the observed heterozygosity of the seedlings was similar between wild populations and specimens seedlings at all sites. At Sites e, g and i, the number of alleles of hypothetical mixed populations were higher than those of the wild populations because one to three unique alleles existed only in the seedlings. The allelic richness of the hypothetical mixed

218 populations was higher than that of the wild populations only at Site e. Although rare alleles were
219 much more common in the wild populations than in the seedlings at all sites, the seedlings had some
220 rare alleles at all sites. Unique alleles in the specimens' seedlings were found at the three sites except
221 Site d. At Sites e and i, the expected heterozygosity of the hypothetical mixed populations was more
222 than that of the wild populations. The inbreeding coefficient of the seedlings ranged from -0.767 to
223 -0.205 , whereas that of the seedlings at Site i indicated significant heterozygosity excess (Table 2).
224 On the other hand, no wild populations exhibited significant heterozygosity excesses. Significant
225 linkage disequilibrium occurred at Site g for only one pair of loci (*Vpy002/Vpy013*; $P < 0.05$).
226
227 Genetic differentiation and structure of herbarium specimens' seedlings and wild populations
228 The STRUCTURE analysis indicates that the wild populations were divided into distinct genetic
229 clusters (Fig. 3a). The ΔK value representing the hierarchical approach for the STRUCTURE
230 analysis was clearly the highest at $K = 2$ (Fig. S1a). In addition, the variance of log likelihood
231 between runs was low and the ΔK value was high at $K = 3$ (Fig. S1a). Therefore, $K = 3$ also yielded
232 meaningful results. Thus, the results obtained with $K = 2$ and $K = 3$ are shown (Fig. 3a). When $K = 2$,
233 the individuals were divided into two clusters. The wild populations at Sites d, e and i were assigned
234 to cluster I, and the wild population at Site g was assigned to cluster II. The F value of cluster I was
235 higher than that of cluster II, and the expected heterozygosity of clusters I and II were 0.780 and
236 0.747, respectively (Fig. 3a). When $K = 3$, cluster I from the $K = 2$ analysis was divided into two
237 clusters. The wild population at Site g was assigned to cluster II and those at Sites e and i were
238 assigned to cluster III. The F value of cluster III was lower than that of cluster I and II, and the
239 expected heterozygosity of cluster I, II and III were 0.743, 0.742 and 0.781, respectively (Fig. 3a).

The wild populations and specimens' seedlings were also divided into distinct genetic clusters (Fig. 3b). The ΔK value representing the hierarchical approach for the STRUCTURE analysis was clearly highest at $K = 3$ (Fig. S1b). Thus, $K = 3$ was the uppermost hierarchical level of the genetic structure. In addition, the variance of log likelihood between runs was low and the ΔK value was high at $K = 4$ (Fig. S1b). Therefore, $K = 4$ also yielded meaningful results. Thus, the results obtained with $K = 3$ and $K = 4$ are shown (Fig. 3b). When $K = 3$, the individuals were clearly divided into three clusters. The wild population and herbarium specimens' seedlings at Site d and the seedlings at Site e were assigned to cluster I. The wild populations at Sites e, g and i were assigned to cluster II. The seedlings at Sites g and i were assigned to cluster III. The F values of cluster I and III were higher than that of cluster II, and the expected heterozygosity of clusters I, II and III were 0.672, 0.787 and 0.584, respectively (Fig. 3b). When $K = 4$, cluster II from the $K = 3$ analysis was divided into two clusters. The wild population at Site g was assigned to cluster II and the wild populations at Sites e and i were assigned to cluster III. The F values of cluster I and IV were higher than those of cluster II and III, and the expected heterozygosity of cluster I, II, III and IV were 0.667, 0.744, 0.786 and 0.584, respectively (Fig. 3b).

According to PCoA based on co-dominant genotypic distances, about 48.9% of the total variation was described by the first two axes (Fig. 4). The specimens' seedlings and the wild individuals were plotted as a single group at Site d. On the other hand, the plots of specimens' seedlings at Sites e and g were nearby the wild individuals. The plots of seedlings at Site i were markedly removed from those of the wild individuals.

Discussion

We found that 18.4% of specimens' seeds were germinated. Furthermore, the specimens' seedlings had unique alleles that did not exist in the wild populations, although the genetic diversity of germinated specimens' seedlings was lower than that of wild populations and clear genetic structure was observed between specimens' seedlings and wild populations at the three sites. These result suggested that viable herbarium-specimen sourced seeds can contribute to the restoration of genetic diversity in threatened plants as *ex situ* conservation resources.

Restoration of genetic diversity by specimens' seedlings

The specimens' seedlings from each site had one to three unique alleles (Table 2). Although we could not estimate the past genetic diversity in each site, the number of alleles of *V. pycnostelma* may have declined in recent decades. The specimens' seedlings probably had the alleles that had extirpated from current wild populations. Thus, the reintroduction of specimens' seedlings with these unique alleles would be useful to restore or augment the genetic diversity of the wild populations.

The genetic diversity (i.e. number of alleles, allelic richness and expected heterozygosity) of specimens' seedlings was lower than that of the wild population. In this study, we used only one specimen fruit per population in order to minimize the damage to the specimens. Because the subfamily Asclepiadoideae of Apocynaceae is pollinated via transfer of pollinia, multiple paternities are very low within fruits (Broyles and Wyatt 1990; Wyatt and Broyles 1994). Therefore, the genetic diversity of the specimens' seedlings would be enhanced by using the seeds of (1) multiple fruits and specimens, (2) species that do not form pollinia (i.e. Orchidaceae and Asclepiadoideae) and allogamous species and (3) fruits from chasmogamous flowers.

Except at Site d, the allele compositions were different and clear genetic structures were

observed between the specimens' seedlings and the wild populations (Figs. 3 and 4). This could be because the specimens' seedlings retained different alleles than the wild populations and they exhibited low genetic diversity. Similarly, significant genetic differences between the seedlings derived from soil seedbanks and the wild populations were also reported (Honnay et al. 2008). Reintroduction of many seedlings taken from only a few specimens or fruits might lead to the dominance of a few alleles in the wild populations. Thus, collecting seeds from multiple fruits and specimens would enhance the genetic diversity of specimens' seedlings so that they more closely resemble the original populations, and the allele frequencies of the seedlings would then be more similar to the wild populations.

Usability of herbarium specimens' seeds of threatened plants

In this study, we determined that 18.4 % of specimens' seeds, which had been collected recently (<19 years), were germinated. These germinated seeds preserved in herbarium could be useful in reintroduction or augmentation resources because viable seeds and bulbils of plants can be preserved for a long time (>20 years) under ideal conditions (Probert et al. 2009; Alsos et al. 2013). In addition, seeds that possess physical dormancy and those with large embryos and little endosperm remain viable for a longer time (Merritt et al. 2014). The seeds of *V. pycnostelma* also possess a physiological dormancy and *Vincetoxicum* species have large embryos (Martin 1946; Zhou et al. 2003; Baskin and Baskin 2014). Thus, the seeds of *V. pycnostelma* specimens collected more than 20 years prior to the study would have a potential to germinate, although we used only nine fruits for germination test in this study. Furthermore, 3.0% of the ungerminated specimens' seeds that were collected 9–18 years prior to the study were viable (Table 1). Because such seeds may be in a

dormant state, seed dormancy should be broken using various methods such as cold stratification, chemicals, heat shock, hormones or scarification (Fontaine et al. 1994; Susko et al. 2001; Kanmegne and Omokolo 2008; Baskin and Baskin 2014). To germinate seeds, it is important to create appropriate environments for inducing germination based on the germination characteristics of the target species. Zygotic embryo culture and callus culture technologies are also available for regenerating endangered plants (Gomes et al. 2003; Rambabu et al. 2006). These technologies could be used to increase the number of individuals derived from the specimens' seeds. The storage conditions in herbaria are also important for specimens' seed longevity. Insect controls that avoid heating, but employ nitrogen, argon, carbon dioxide and freezing have been developed in herbaria (Strang 1992; Valentin 1993); these controls do not have negative effects on seed germination of many species (Bass and Stanwood 1978; Prokopiev et al. 2014).

It is recommended that sampling regimes for reintroduction resources should involve collecting 1 to 20 seeds per individual from each of 10 to 50 individuals belonging to each of five separate populations to remove the inbreeding depression (Falk and Holsinger 1991). Hoban and Schlarbaum (2014) also suggest that we should collect seeds from 25 to 30 individuals belonging to the few but widely dispersed populations and between two to eight seeds of fruits per individual to maintain their genetic diversity. However, many specimens of target species collected from the same site (>10 individuals) may be rarely preserved in a herbarium. In addition, damage to an excessive number of specimens should be minimized because museum specimens are very valuable and irreplaceable (Graves and Braun 1992; Wandeler et al. 2007; Shiga 2013). Furthermore, the number of retained and available seeds on herbarium specimens would be also not known before their germination tests. Thus, it may be difficult to collect sufficient seeds from specimens in only a few herbaria as

reintroduction resources. For example, it is feasible to sample across many sites and have wider genetic variation if material is sourced from several herbaria.

In recent decades, in addition to the designation of the sampling guidelines of seed banking projects, their utilization for *ex situ* conservation has been increasingly applied to a more diverse array of wild species (Schoen and Brown 2001; Van Slageren 2003; Guerrant et al. 2014; Hoban and Schlarbaum 2014). However, the application of seed banking projects may be difficult for the conservation of locally threatened populations because these projects usually do not assume the conservation of the local populations. The use of herbaria specimens' seeds would also remarkably contribute to restore the genetic diversity of not only the plant species but also each locally threatened population if the viable specimens' seeds collected at the target species or populations are preserved in herbaria.

Acknowledgments

We thank M. Azuma and M. Zushi for their helpful specimens' seeds sampling support. This work was supported by Grants in Aid for Scientific Research (23701024, 26350387) from the Japan Society for the Promotion of Science.

350 **References**

- 351 Aguilar R, Ashworth L, Galetto L, Aizen MA (2006) Plant reproductive susceptibility to habitat
352 fragmentation: review and synthesis through a meta-analysis. *Ecol lett* 9: 968–980
- 353 Alsos IG, Müller E, Eidesen PB (2013) Germinating seeds or bulbils in 87 of 113 tested Arctic species
354 indicate potential for ex situ seed bank storage. *Pol Biol* 36: 819–830
- 355 Baskin CC, Baskin JM (2014) *Seeds: Ecology, Biogeography, and Evolution of Dormancy and*
356 *Germination* (Second Edition). Academic Press, San Diego
- 357 Bass LN, Stanwood PC (1978) Long-term preservation of sorghum seed as affected by seed moisture,
358 temperature, and atmospheric environment. *Crop Sci* 18: 575–577
- 359 Becker U, Colling G, Dostal P, Jakobsson A, Matthies D (2006) Local adaptation in the monocarpic
360 perennial *Carlina vulgaris* at different spatial scales across Europe. *Oecologia* 150: 506–518
- 361 Bewley JD, Bradford KJ, Hilhorst HWM, Nonogaki H (2013) *Seeds: Physiology of Development,*
362 *Germination and Dormancy*, 3rd Edition. Springer, New York
- 363 Blows MW, Hoffmann AA (2005) A reassessment of genetic limits to evolutionary change. *Ecology* 86:
364 1371–1384
- 365 Boutin–Ganache I, Raposo M, Raymond M, Deschepper CF (2001) M13–tailed primers improve the
366 reliability and usability of microsatellite analyses performed with two different allele sizing methods.
367 *Biotechniques* 31: 24–26
- 368 Bowles ML, Betz RF, DeMauro MM (1993) Propagation of rare plants from historic seed collections:
369 implications for species restoration and herbarium management. *Restor Ecol* 1: 101–106
- 370 Broyles SB, Wyatt R (1990) Paternity analysis in a natural population of *Asclepias exaltata*: multiple
371 paternity, functional gender, and the "pollen-donation hypothesis". *Evolution* 44: 1454–1468

- 372 Ceballos G, Ehrlich PR (2002) Mammal population losses and the extinction crisis. *Science* 296: 904–907
- 373 Cottrell, HJ (1947) Tetrazolium salt as a seed germination indicator. *Nature* 159: 748
- 374 Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing
375 STRUCTURE output and implementing the Evanno method. *Conserv Genet Res* 4: 359–361
- 376 Elias SG, Copeland LO, McDonald MB, Baalbaki RZ (2012) Seed Testing: Principles and Practices.
377 Michigan State University Press, East Lansing
- 378 El Mousadik EA, Petit RJ (1996) High level of genetic differentiation for allelic richness among
379 populations of the argan tree [*Argania spinosa* (L.) Skeels] endemic to Morocco. *Theor Appl Genet* 92:
380 832–839
- 381 Environment Agency of Japan (2000) Threatened Wildlife of Japan – Red Data Book 2nd ed. Vascular 3.
382 Plants. Japan Wildlife Research Center, Tokyo (in Japanese)
- 383 Falk DA, Holsinger KE (1991) Genetics and conservation of rare plants. Oxford university press, New
384 York
- 385 Flora of Korea Editorial Committee (2007) The Genera of Vascular Plants of Korea. Academy Publishing,
386 Seoul
- 387 Fontaine O, Huault C, Pavis N, Billard JP (1994) Dormancy breakate of *Hordeum vulgare* seeds: effects
388 of hydrogen peroxide and scarification on glutathione level and glutathione reductase activity. *Plant*
389 *Physiol Biochem* 32: 677–683
- 390 Frankel OH, Soulé ME (1981) Conservation and Evolution. Cambridge University Press, Cambridge
- 391 Frankham R, Ballou JD, Briscoe DA (2009) Genetic management of captive populations. In: Introduction
392 To Conservation Genetics, Second Edition. Cambridge university press, Cambridge, pp 430–452
- 393 Gomes GAC, Paiva R, de Oliveira PPD, de Santiago EJA (2003) Plant regeneration from callus cultures

- 394 of *Maclura tinctoria*, an endangered woody species. In Vitro Cell Dev–Pl 39: 293–295
- 395 Gottelli D, Sillero–Zubiri C, Appelbaum GD, Roy MS, Girman DJ, Garcia–Moreno J, Ostrander EA,
396 Wayne RK (1994) Molecular genetics of the most endangered canid: the Ethiopian wolf *Canis simensis*.
397 Mol Ecol 3: 301–312
- 398 Goudet J (2001) FSTAT, a program to estimate and test gene diversities and fixation indices (version
399 2.9.3). <http://www2.unil.ch/popgen/softwares/fstat.htm>. Accessed 14 Jan 2014
- 400 Graves GR, Braun MJ (1992) Museums: storehouses of DNA? Science 255: 1335–1336
- 401 Guerrant Jr EO, Havens, K, Vitt P (2014) Sampling for effective ex situ plant conservation. Int J Plant Sci
402 175: 11–20
- 403 Hereford J (2009) A quantitative survey of local adaptation and fitness trade-offs. Amer Nat 173: 579–588
- 404 Hoban S, Schlarbaum S (2014) Optimal sampling of seeds from plant populations for ex-situ
405 conservation of genetic biodiversity, considering realistic population structure. Biol Conserv 177: 90–
406 99
- 407 Honnay O, Bossuyt B, Jacquemyn H, Shimono A, Uchiyama K (2008) Can a seed bank maintain the
408 genetic variation in the above ground plant population? Oikos 117: 1–5
- 409 Hooper DU, Chapin Iii FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau M,
410 Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle DA (2005) Effects of biodiversity
411 on ecosystem functioning: a consensus of current knowledge. Ecol monographs 75: 3–35
- 412 Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009) Inferring weak population structure with the
413 assistance of sample group information. Mol Ecol Resour 9: 1322–1332
- 414 Huff DD, Miller LM, Chizinski CJ, Vondracek B (2011) Mixed-source reintroductions lead to
415 outbreeding depression in second–generation descendants of a native North American fish. Mol ecol

- 416 20: 4246–4258
- 417 Kanmegne G, Omokolo ND (2008) Germination of *Garcinia kola* (Heckel) seeds in response to different
- 418 hormone treatments. *Fruits* 63: 155–161
- 419 Keller LF, Waller DM (2002) Inbreeding effects in wild populations. *Trends Ecol Evolut* 17: 230–241
- 420 Koyanagi TF, Furukawa T (2013) Nation-wide agrarian depopulation threatens semi-natural grassland
- 421 species in Japan: Sub-national application of the Red List Index. *Biol Conserv* 167: 1–8
- 422 Leimu R, Fischer M (2008) A meta-analysis of local adaptation in plants. *PLoS One* 3: e4010
- 423 León-Lobos P, Way M, Aranda PD, Lima-Junior M (2012) The role of ex situ seed banks in the
- 424 conservation of plant diversity and in ecological restoration in Latin America. *Plant Ecol Divers* 5:
- 425 245–258
- 426 Lledó MD, Crespo MB, Amo-Marco JB (1996) Micropropagation of *Limonium thiniense* Erben
- 427 (Plumbaginaceae) using herbarium material. *Bot Gard Micropropag News* 2: 18–21
- 428 Magrini S (2011) Herbaria as useful spore banks for integrated conservation strategies of pteridophytic
- 429 diversity. *Plant Biosyst* 145: 635–637
- 430 Magrini S, Olmati C, Onofri S, Scoppola A (2010) Recovery of viable germplasm from herbarium
- 431 specimens of *Osmunda regalis* L. *Amer. Fern J.* 100: 159–166
- 432 Martin AC (1946) The comparative internal morphology of seeds. *Amer Midl Nat* 36: 513–660
- 433 Mattila AL, Duploux A, Kirjokangas M, Lehtonen R, Rastas P, Hanski I (2012) High genetic load in an
- 434 old isolated butterfly population. *Proc Natl Acad Sci* 109: 2496–2505
- 435 Merritt DJ, Martyn AJ, Ainsley P, Young RE, Seed LU, Thorpe M, Hay FR, Commander LE, Shackelford
- 436 N, Offord CA, Dixon KW, Probert RJ (2014) A continental-scale study of seed lifespan in
- 437 experimental storage examining seed, plant, and environmental traits associated with longevity.

- 438 Biodivers Conserv 23: 1081–1104
- 439 Milián-García Y, Venegas-Anaya M, Frias-Soler R, Crawford AJ, Ramos-Targarona R,
440 Rodríguez-Soberón R, Alonso-Tabet A, Thorbjarnarson J, Sanjur OI, Espinosa-López G, Bermingham
441 E (2011) Evolutionary history of Cuban crocodiles *Crocodylus rhombifer* and *Crocodylus acutus*
442 inferred from multilocus markers. J Exp Zool A Ecol Genet Physiol, 315: 358–375
- 443 Milligan B (1992) Plant DNA isolation. In: Hoelzel AR (ed) Molecular Genetic Analysis of Populations: a
444 Practical Approach. IRL Press, Oxford, pp 59–88
- 445 Ministry of the Environment Government of Japan (2012) The Red List of Vascular Plants of Japan.
446 https://www.env.go.jp/press/file_view.php?serial=20557&hou_id=15619. Accessed 14 Jan 2014
- 447 Montalvo AM, Ellstrand NC (2001) Nonlocal transplantation and outbreeding depression in the subshrub
448 *Lotus scoparius* (Fabaceae). Am J of Bot 88: 258–269
- 449 Nakahama N, Kaneko S, Hayano A, Isagi Y, Inoue–Murayama M, Tominaga T (2012) Development of
450 microsatellite markers for the endangered grassland species *Vincetoxicum pycnostelma* (Apocynaceae),
451 using next–generation sequencing technology. Conserv Genet Resour 4: 669–671
- 452 Osaka Museum of Natural History (2012) Collecting and storage business. Annu Rep Osaka Mus Nat
453 Hist 37: 19–27 (in Japanese)
- 454 Palomares F, Godoy JA, López–bao JV, Rodríguez A, Roques S, Casas–Marce M, Revilla E, Delibes, M
455 (2012) Possible extinction vortex for a population of Iberian lynx on the verge of extirpation. Conserv
456 Biol 26: 689–697
- 457 Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for
458 teaching and research. Mol Ecol Notes 6: 288–295
- 459 Pritchard JK, Wen X, Falshu D (2010) Documentation for STRUCTURE ver. 2.3. University of Chicago,

- 460 Chicago
- 461 Probert RJ, Daws MI, Hay FR (2009) Ecological correlates of ex situ seed longevity: a comparative study
462 on 195 species. *Ann Bot* 104: 57–69
- 463 Prokopiev IA, Filippova GV, Shein AA (2014) Effect of different conditions of welsh onion seed storage
464 on germination and cytogenic characteristics of its seedlings. *Russ J Genet* 4: 614–617
- 465 Qvenild M (2008) Svalbard Global Seed Vault: a ‘Noah’s Ark’ for the world’s seeds. *Dev Pract* 18: 110–
466 116
- 467 Rambabu M, Upender M, Ujjwala D, Ugandhar T, Praveen M, Swamy NR (2006) In vitro zygotic
468 embryo culture of an endangered forest tree *Givotia rottleriformis* and factors affecting its germination
469 and seedling growth. *In Vitro Cell Dev–Pl* 42: 418–421
- 470 Schoen DJ, Brown AH (2001) The conservation of wild plant species in seed banks. Attention to both
471 taxonomic coverage and population biology will improve the role of seed banks as conservation tools.
472 *Bioscience* 51: 960–966
- 473 Shiga T (2013) A relationships among curator, collector, and user surrounding natural history
474 specimens: toward a better specimen preservation, collection and usage (*Museum and ecology* (22)).
475 *Jap J Ecol* 63: 375–383 (in Japanese)
- 476 Smouse PE, Peakall R (1999) Spatial autocorrelation analysis of individual multiallele and multilocus
477 genetic structure. *Heredity* 82: 561–573
- 478 Strang TJK (1992) A review of published temperatures for the control of pest insects in museums. *Collect*
479 *Forum* 8: 41–67
- 480 Susko DJ, Mueller, JP, Spears, JF (2001) An evaluation of methods for breaking seed dormancy in kudzu
481 (*Pueraria lobata*). *Can J Bot* 79: 197–203

- 482 Uematsu Y, Koga T, Mitsuhashi H, Ushimaru A (2010) Abandonment and intensified use of agricultural
483 land decrease habitats of rare herbs in semi-natural grasslands. *Agricult Ecosys Environ* 135: 304–309
- 484 Uesugi R, Nishihiro J, Tsumura Y, Washitani I (2007) Restoration of genetic diversity from soil seed
485 banks in a threatened aquatic plant, *Nymphoides peltata*. *Conserv Genet* 8: 111–121
- 486 Valentin N (1993) Comparative analysis of insect control by nitrogen, argon and carbon dioxide in
487 museum, archive and herbarium collections. *Int Biodeterior Biodegrad* 32: 263–278
- 488 Van Slageren W (2003) The Millennium Seed Bank: building partnerships in arid regions for the
489 conservation of wild species. *J Arid Environ* 54: 195–201
- 490 Washitani I (1987) A convenient screening test system and a model for thermal germination responses of
491 wild plant seeds: behavior of model and real seeds in the system. *Plant Cell Environ* 10: 587–598
- 492 Wandeler P, Paquita EAB, Lukas FK (2007) Back to the future: museum specimens in population genetics.
493 *TREE* 22: 634–642
- 494 Willi Y, Van Buskirk J, Hoffmann AA (2006) Limits to the adaptive potential of small populations. *Annu*
495 *Rev Ecol Evol Syst* 37: 433–458
- 496 Windham MD, Wolf PG, Ranker TA (1986) Factors affecting prolonged spore viability in herbarium
497 collections of three species of *Pellaea*. *Amer Fern J* 76: 141–148
- 498 Wu ZY, Raven PH (1995) *Flora of China*. Vol. 16 (Gentianaceae through Boraginaceae). Science Press,
499 Beijing, and Missouri Botanical Garden Press, St. Louis
- 500 Wyatt R, Broyles S (1994) Ecology and evolution of reproduction in Milkweeds. *Annu Rev Ecol Evol* 25:
501 423–441
- 502 Yamazaki T (1993) *Asclepiadaceae*. In: Iwatsuki K, Yamazaki T, Boufford DE, Ohba H. (Eds), *Flora of*
503 *Japan IIIa*. Kodansha, Tokyo, pp 176 (in Japanese)

- 504 Zaghoul M, Reisch C, Poschlod P (2013) Soil seed bank contributes significantly to genetic variation of
- 505 *Hypericum sinaicum* in a changing environment. Plant Syst Evol 299: 819–828
- 506 Zhou YF, Hang YZ, Zhou LL, Wang KC (2003) The germination characteristics of *Cynanchum*
- 507 *paniculatum* seeds. J Plant Resourc Environ 12: 35–38
- 508
- 509
- 510
- 511
- 512
- 513
- 514
- 515
- 516
- 517
- 518
- 519
- 520
- 521
- 522
- 523
- 524
- 525

526 **Supplementary material**

527 Additional supplemental material can be found in the online version of this article.

528

529 **Table S1** The condition of germination test

530

531 **Figure S1** The ΔK in the STRUCTRE analysis

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548 **Table 1** Specimen characteristics and number of seeds used in experiments

Site	Prefecture	No. of years after collected	Date of collecting specimens	Voucher specimen	Increasing temperature condition			decreasing temperature condition		
					No. of seeds used experiment	No. of germinated seeds	No. of viable ungerminated seeds	No. of seeds used experiment	No. of germinated seeds	No. of viable ungerminated seeds
a	Gifu	16	25 Sep. 1997	H. Marui 2434 (OSA203391)	10	0	1	10	0	0
b	Kyoto	51	16 Sep. 1962	M. Hutoh 24248 (OSA214994)	9	0	0	9	0	0
c	Kyoto	22	24 Sep. 1991	T. Fujii 2413 (OSA209484)	10	0	0	10	0	0
d	Nara	4	12 Oct. 2009	S. Onoue s.n. (OSA225354)	9	3	0	9	3	0
e	Nara	16	03 Aug. 1997	K. Seto 47673 (OSA104247)	11	0	0	11	4	0
f	Nara	62	03 Jul. 1951	M. Hori s.n. (OSA190777)	8	0	0	8	0	0
g	Osaka	3	26 Nov. 2010	F. Uwakubo 101126-002 (OSA185864)	12	0	0	12	2	0
h	Osaka	9	10 Oct. 2004	K. Hirano 2501 (OSA280257)	11	0	2	11	0	0
i	Hyogo	18	02 Nov. 1994	S. Miyake 2987 (OSA122127)	23	16	2	23	10	0
Total					103	19	5	103	19	0

549

550

Table 2 Genetic diversity measurements of each site of *Vincetoxicum pycnostelma*. Each site contained specimens' seedlings, a wild and a hypothetical mixed population (specimens' seedlings + wild population). *A*, numbers of alleles per locus; *AR*, allelic richness; *RA*, summed number of rare alleles; *UA*, summed number of unique alleles that are only present in specimens' seedlings; *H_O*, observed heterozygosity (in bold numbers, if values are significantly deviated from HWE); *H_E*, expected heterozygosity; *F_{IS}*, inbreeding coefficient

Site	Wild Population size	Sample	No. of samples	<i>A</i>	<i>AR</i>	<i>RA</i>	<i>UA</i>	<i>H_O</i>	<i>H_E</i>	<i>F_{IS}</i>
d	20	specimens' seedlings	6	3.111	2.366	5	0	0.815	0.582	-0.382
		Wild population	19	6.222	2.683	17		0.754	0.687	-0.106
		Hypothetical mixed population	25	6.222	2.637	17		0.769	0.680	-0.137
e	220*	specimens' seedlings	4	3.000	2.406	5	1	0.778	0.566	-0.348
		Wild population	46*	11.333	2.917	51		0.749	0.752	0.008
		Hypothetical mixed population	50	11.444	2.929	52		0.751	0.756	0.010
g	100	specimens' seedlings	2	2.222	2.222	2	1	0.833	0.472	-0.767
		Wild population	50	10.000	2.859	39		0.767	0.748	-0.030
		Hypothetical mixed population	52	10.111	2.858	40		0.769	0.748	-0.034
i	24	specimens' seedlings	20	2.889	2.230	3	3	0.694	0.570	-0.205
		Wild population	16	7.556	2.826	25		0.701	0.723	0.030
		Hypothetical mixed population	36	8.000	2.765	27		0.698	0.725	0.044

*: sampled about 1 km from where specimens were originally collected, because original population was locally extinct

Figure captions

Fig. 1 a The herbarium specimen of *Vincetoxicum pycnostelma* collected site i (OSA122127). Arrow head indicates the fruit, Bar indicates 50mm. **b** Seeds in the fruits of OSA122127. Bar indicates 5.0mm. **c** Habitat of *V. pycnostelma* (Site d). **d** Juvenile individuals germinated from the seeds of herbarium specimens (right pot: OSA104247, left pot: OSA185864)

Fig. 2 a Locations of the collected *Vincetoxicum pycnostelma* specimens in Kinki and Tokai districts. **b** Location of Kinki and Tokai districts in Japan

Fig. 3 Results of Bayesian clustering in the STRUCTURE analysis (Pritchard et al. 2010). **a** The proportion of the membership coefficient of 131 individuals in the wild populations at the four sites for each of the inferred clusters for $K = 2$. **b** The proportion of the membership coefficient of 163 individuals in the specimens' seedlings and wild populations at the four sites (right side: specimens' seedlings, left side: wild populations) for each of the inferred clusters for $K = 3$ and $K = 4$. Each column represents an individual

Fig. 4 Principal coordinates analysis plots of individuals at the four sites (each contains specimens' seedlings and wild populations) based on co-dominant genotypic distances (Smouse and Peakall 1999). Red, green, white and blue symbols represent Sites d, e, g and i, respectively. Axis 1 explains 27.5% of the variance and axis 2 explains 21.4% of the variance

Figure 1




Figure 2




Figure 3





Figure 4




1 **Table S1** Relationships between temperature and number of days in each condition. In the increasing temperature condition, the temperature was
2 alternated between 12 and 25 °C in 24-h intervals after arriving at the final temperature

Increasing temperature condition	Temperature (°C)	4	8	12	16	20	24	28	32	36	12~25
	Number of days	8	5	4	3	2	2	2	2	2	5
Decreasing temperature condition	Temperature (°C)	36	32	28	24	20	16	12	8	4	25
	Number of days	2	2	2	2	2	3	4	5	8	5

3
4
5
6
7

Figure S1

The ΔK in the STRUCTRE analysis based on the rate of change in the log probability of data between successive K values (Evanno et al. 2005). (a) The result of 131 individuals in the only wild populations of the four sites. (b) The result of 163 individuals in the specimens' seedlings and wild populations of four sites.



References

Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software Structure: a simulation study. Mol Ecol 14: 2611–2620